Sedation/Tranquilization, Anesthesia and

MOUSE FORMULARY

DRUG NAME and DOSE	ROUTE& FREQUENCY	NOTES
Anesthetic Induction		
Sodium Pentobarbital (Nembutal) 30-90mg/kg	IP	Beware of dose related deep anesthetic plane, respiratory depression, narrow margin of safety, hypothermia and prolonged recovery (ref # 1 & 11)
Ketamine 100mg/kg + Xylazine 10mg/kg	IP	General anesthesia for surg J L F D O S U R F H G X U H V ' R Q R W V X S S O H P H Q W
Ketamine 100mg/kg + Diazepam OR Midazolam 5mg/kg	IP	General anesthesia for surg L F D O S U R F H G X U I ' R Q R W V X S S O H P H Q W
Ketamine 50-75mg/kg +		

		hypothermia and prolonged recovery (ref # 1 & 11)
Ketamine 40-80mg/kg + Xylazine 5-10mg/kg	IP	General anesthesia for surg L F D O S U R F H G X U H V Q R W V X S S O H P H Q W
Ketamine 75mg/kg + Diazepam OR Midazolam 5mg/kg	IP	
Ketamine 50-75mg/kg + Dexmedetomidine 0.25mg/kg	IP	
Isoflurane	1-4%	Generally first choice agent in rodents because it c be easily titrated to deliver dose required for anesthesia Induce rodent in a chamber at 3-4%, then reduce to 1-2% for procedure
Analgesics		
Meloxicam 1-2mg/kg	PO, SQ q12	

	Mole Rat	PO, SC; SID	
	OPIOIDS		
Buprenorphine	Rodents	SC, PO; QID-BID	
Morphine	Naked Mole Rat		AVOID – causes hyperalgesia and severe aggression

RABBIT FORMULARY

DRUG NAME and DOSE	ROUTE & FREQUENCY	NOTES
Sedation		
Acepromazine 0.75mg/kg	IM	Produces a peripheral vasodilation useful for venipuncture
Dexmedetomidine 0.05mg/kg	SQ	Produces moderate sedation, useful for minimally invasive procedures less than 30 minutes
Induction		
Ketamine 10-20mg/kg + Dexmedetomidine 0.1mg/kg	SQ	