Hospice & Palliative Care Association of New York State

Palliative Care Information Act Resource

Table of Contents

I. What is Palliative Care & How is it different from Hospice?

a) Palliative Care Defined

b) Hospice Defined

c) How is Palliative Care Different from Hospice?

II. Appropriateness

a) Who needs Palliative Care? How do I know this person is appropriate for Palliative Care?

b) What is the benefit of Palliative Care?

i. For the patient & family

ii. For the provider

III. Timing

a) When is a Hospice referral appropriate based on diagnosis?

IV. Communication

a) How do I begin the conversation?

b) How do I engage the family in the conversation?

c) Cultural considerations for communication

d) Spiritual considerations for communication

V. Access

a) How is Palliative Care paid for?

b) How do I find a Palliative Care Provider?

VI. Role

a) What is the continuing role of the Primary Care Provider when a Palliative Care referral is made?

VII. Additional Resources

I.

Question:

the family; the Hospice team works with the patient and family to develop a personalized plan of care that incorporates what is important to the patient and the caregiver

The following graph provides a v

The findings of a recent study called '*Early Palliative Care for Patients with Metastatic Non–Small-Cell Lung Cancer*' which was reported in the New England Journal of Medicine Article state that compared with patients receiving standard care, patients who received palliative care early after diagn-3.g(s)-5its

nontonday2 dy2 04( )]TJ -0.004 Tc 0.004 Tw [experienedssiant

Predicting a life-expectancy of 6 months or less usually involves a significant, documented deterioration in physical status/function such as weight loss or decreased function and/or an end stage disease. Decline may also be due to refusal of treatments, medications or hospitalization aimed at improving or stabilizing an advanced disease.
A patient can be appropriate for hospice even without a specific end-stage disease (*see DECLINE IN STATUS*).

- AMYOTROPHIC LATERAL SCLEROSIS (ALS): Patients tend to have a constant overall rate of decline, whether rapid or slow. They should have 1 OR 2 OR 3 below:

1. CRITICALLY IMPAIRED BREATHING CAPACITY occurring over the past 12 months with ALL of the following: a) dyspnea at rest; b) supplemental O2 at rest; c) no ventilator or ventilator only for comfort. A vital capacity <30% is supportive, if available OR
2. RAPID PROGRESSION AND CRITICAL NUTRITIONAL IMPAIRMENT with ALL of the following in the past 12 months: a) bed-bound; b) barely or unintelligible speech; c) pureed diet; d) major assist in all ADLs; e) insufficient oral intake; f) continuing weight loss; g) NO

1. PT > 5 SEC. OVER CONTROL OR INR. 1.5 AND Albumin < 2.5 AND
2. ONE OF THE FOLLOWING: a) refractory ascites or non-compliance; OR b) spontaneous bacterial peritonitis; OR c) hepatorenal syndrome; OR d) refractory encephalopathy or non-compliance; OR e) recurrent esophageal variceal bleeding despite treatment
3. SUPPORTIVE: a) weight loss>10%; b) muscle wasting/loss of strength; c) continued alcohol consumption; d) hepatocellular cancer; e) HBsAg positive; or f) hepatitis C refractory to treatment

- LUNG DISEASE: patients should have 1 AND 2; factors in 3 are supportive

1. Dyspnea at rest and minimal exercise tolerance (with FEV1 <30% IF AVAILABLE) and progression of disease with increased visits, hospitalizations or MD/ER visits (documented serial decrease in FEV1 40 ml/year if available) AND
2. pO2 <55 mmHg ON ROOM AIR OR O2 SAT < 88 ON O2 OR Pco2 >50 mm Hg
3. SUPPORTIVE: a) cor pulmonale; b) weight loss.10% in past 6 months; c) HR > 100/min at rest

- KIDNEY DISEASE: patients should have 1 AND EITHER 2 OR 3; factors in 4 and 5 are supportive

1. Not seeking dialysis or transplant or stopping dialysis AND
2. Creatinine clearance < 10 cc/min (<15 FOR DIABETICS) OR LESS THAN 15 cc/min WITH CHF (< 20 cc/min FOR DIABETICS) OR
3. CREATININE > 8 mg/dl (> 6 FOR DIABETICS)
4. SUPPORTIVE FOR ACUTE RENAL FAILURE: a) mechanical ventilation; b) cancer; c) chronic lung disease; d) advanced heart disease; e) advanced liver disease; f) sepsis; g) AIDS; h)

C. Signs: 1) systolic BP<90 or postural hypotension; 2) ascites; 3) venous, arterial or lymphatic obstruction; 4) edema; 5) pleural/ pericardial effusion; 6) weakness; 7) change in LOC
D. Laboratory (NOT required): 1) increasing CO2 or decreasing pO2 or O@ sat; 2)

underscore, and these can be powerful motivators. By preparing for the worst—by making a living will, naming a health care proxy, preparing financial matters, or settling family affairs—patients can address fears, clarify priorities, and strengthen relationships with loved ones, all components of a good death identified in empirical studies of patients with life-threatening illnesses. Furthermore, some patients may be thinking about the worst but are afraid to discuss it for fear of frightening their family or disappointing their physician. Naming and discussing these concerns, which often

Patient Says:	Physician Responds:
I hope that the treatment will help me out.	I'm also hoping that the treatment will help you. If it works, what will be the most important thing for you to do? I wonder, also, if you would be willing to talk about what we should do in case the treatment doesn't work.
I'm concerned that talking about what might happen will be overwhelming.	Well, talking about what might happen can be frightening at first, but most patients I have worked with have found it helpful in the long run.
I'm worried that talking about what happens if' would be giving up.	I understand your concern, but we are not talking about giving up
I don't want to talk about what I'm worried about.	Okay, I realize that talking about worries can be a hard thing to do. If you do want to talk about these issues in the future, I would be open to it.

I hope that

F = FEELINGS related to the illness, especially fears

- What are you most concerned about?
- Do you have any specific fears or worries right now?
- I imagine you have had many different feelings as you have coped with this illness.
- Sometimes people have fears that they keep to themselves and don't tell their doctor.

I = IDEAS and explanations of the cause of a symptom or illness

- What do you think might be going on?
- What do you think this pain means?
- Do you have ideas about what might have caused this illness?

F = FUNCTIONING , the illness' impact on daily life

- How has your illness affected you day to day?
- What have you had to give up because of your illness?
- What goals do you have now in your life? How has your illness affected your goals?
- How does this illness affect important people in your life?

E = EXPECTATIONS of the doctor & the illness

- What do you expect or hope I can do for you today?
- Do you have expectations about how doctors can help?
- What do you hope this treatment will do for you?
- What are your expectations about what might happen with this illness?

Reference: End of Life/Palliative Education Resource Center

Talking About Treatment Options and Palliative Care: A Guide for Clinicians At Time of Diagnosis: Begin to Understand Your Patient's Priorities and Goals

When patients first learn that they have a potentially life-limiting illness their focus is on curing the disease. However, there are opportunities as an initial plan of care is developed to discuss the benefits of advance care planning and the difference between curative and palliative care. *Introducing the concept of palliative care as care focused on alleviating symptoms even as you work to cure the illness will allow you to continue referencing palliation of symptoms, optimizing functional capacities and the importance of psychosocial support for the patient and the family.*

The questions in this document, when introduced during a conversation about treatment options can provide clinicians with information about what is important to the patient. Simply asking one or two of these questions can lead to future discussions about palliative care and present the patient with an opportunity to explore the topic with you when they are ready.

As we think about treatment options, what is one thing about you that you think is important to share with me?

- What is your biggest fear when you think about your diagnosis?
- What is your biggest fear about the treatments we have discussed?
- What does quality of life mean to you?
- As things progress, how much do you want to know about your disease and all the

When Treatment May Not Be Working: Exploring End-of-Life Options

At some point, it may become apparent that continued treatments are not likely to offer any specific therapeutic benefit for your patient. One question that is often used to frame discussion about ending curative treatment protocols is “Would you be surprised if your patient died in

Understanding Your Patient's Needs: Responding to Patient/Caregiver Inquiries
Listening to what your patient is saying or not saying is a crucial part to understanding their needs. Questions a patient may ask you are listed below, along with what they might be thinking and truly wanting to know.

When a Patient/Caregiver Asks…	They Might be Thinking…
What are my chances?	Realistically, how likely is it that this will work?
What would you do if you were me?	I am overwhelmed, and I need help or guidance in making these decisions.
How long do I have?	I feel like I am dying… *Or* I am scared about facing death, but I don't have the words to express this… *Or* I really want to know what a realistic time line is so I can make good choices and get things in order…
What are the options?	This isn't working. I want to try something else.

Reference: CaringInfo.org

Return to Top

Question:

A cornerstone procedure in Palliative Care is leadership of family meetings to establish goals of care, typically completed at a time of patient change in status, where the value of current treatments needs to be re-evaluated. As with any procedure, preparation is essential to ensure the best outcome.

Following is a framework overview for 'Preparing for the Family Meeting.'
Data Review: Review the medical history relevant to the current medical situation (e.g. history of disease progression, symptom burden, past treatments, treatment-related toxicity, and prognosis).

- Review all current treatments (e.g. renal dialysis, artificial nutrition, antibiotics) and any positive and/or negative treatment effects.
- Review all treatment options being proposed
- Determine the prognosis with and without continued disease-directed treatments. Prognostic information includes data concerning future patient function (physical/cognitive), symptom burden, and time (longevity).
- Solicit and coordinate medical opinions about the utility of current treatments among consultants and the primary physician. If possible, families need to hear a single medical consensus – all relevant clinicians should be contacted and consensus reached prior to the

4. Using the 'D word' if relevant: when a patient is clearly deteriorating and death is likely within the next days to weeks, or even a few months, it is appropriate to use the word *dying* in the conversation. Both patients and surrogates find that using the word *dying*, if done compassionately, is helpful in clearing what is often a confusing and frightening situation. *I'm afraid we have run out of options to shrink the cancer. Based on your declining function, I believe you are dying.*

5. Silence Whether or not you use the word *dying*, when you have presented bad news (such as information about disease progression), the next step is for you to allow silence, and let the family/patient respond. In truth, no matter what you might imagine the response from the patient/family to be once the bad news is delivered, you really cannot predict their emotional reaction (e.g. relief, anxiety, anger, regret, fear, etc). This silence can be uncomfortable; resist the urge to fill it with more facts as they will not be heard. Not all patients/families express emotions at this point and instead respond practically (*What happens next then?*). This is fine, but you need to wait, silently, to see what response the patient/family demonstrates. In addition, even practical questions have underlying emotions (*Are you sure?* Or – *There must be something you can do?*). It is important to respond to both the factual aspect of the question (*Yes I am sure.* Or – *There are no more effective treatments available*.), as well as the emotional level (*I wish I had better news for you.* Or – *I wish our treatments worked better than they do.).*

When the patient/surrogates openly acknowledge that current treatments are no longer effective, that death is coming, they will generally ask one of the following questions: *How long? What will happen? Will there be suffering? What do we do now?* Your response at this point should be to address prognosis in terms of time, function, and symptoms, as best you can.
Authors: David E Weissman MD, Timothy Quill MD, and Robert M Arnold MD
Reference End of Life / Palliative Education Resource Center

Following is a concise overview of running a family meeting:
Moderating a Family Conference ~ Family Conference Process Steps

1. Why are you meeting? Clarify conference goals in your own mind. What do you hope to accomplish?
2. Where: A room with comfort, privacy and circular seating.
3. Who: Patient (if capable to participating)2(aP6)6(tin)2(M)

- Defer discussion of decision making until the next step.
- Respond to emotional reactions (See Tool Facts #29, 59).

7. Family Discussion with a Decisional Patient
   - Ask the patient, "What decision(s) are you considering?"
   -

U – Unique cultural values. Use respectful, curious, and open-ended questions about a

and biases that are derived from their own cultural origin and from the biomedical world view of their professional training. Medicine itself is a cultural system with its own specific language, values, and practices that must be translated, interpreted, and negotiated with patients and their families. This article describes two cases that raise issues about cross-cultural end-of-life practice and suggests strategies for negotiating common problems.

Reference: Strategies for Culturally Effective End-of-Life Care
American College of Physicians-American Society of Internal Medicine 2002, Vol. 136 (9), pg. 673-679

Return to Top

Question:

- As we plan for your medical care near the end of life, in what ways will your religion and spirituality influence your decisions?

Reference: End of Life / Palliative Education Resource Center

Religious traditions and their impact on healthcare decision-making. This resource for clinicians includes brief articles on common ways in which spiritual considerations might affect health care decisions