

Bush-Francis Catatonia Rating Scale

1. Excitement:

Extreme hyperactivity, constant motor unrest which is apparently non-purposeful. Not to be attributed to akathisia or goal-directed agitation.

- 0= Absent
- 1= Excessive motion, intermittent.
- 2= Constant motion, hyperkinetic without rest periods.
- 3= Full-blown catatonic excitement, endless frenzied motor activity.

2. Immobility/ Stupor:

Extreme hypoactivity, immobile, minimally responsive to stimuli

- 0= Absent
- 1= Sits abnormally still, may interact briefly.
- 2= Virtually no interaction with external world.
- 3= Stuporous, non-reactive to painful stimuli.

3. Mutism:

Verbally unresponsive or minimally responsive.

- 0= Absent
- 1= Verbally unresponsive to majority of questions; incomprehensible whisper.
- 2= Speaks less than 20 words/5 minutes.
- 3= No speech.

4. Staring:

Fixed gaze, little or no visual scanning of environment, decreased blinking.

- 0= Absent
- 1= Poor eye contact, repeatedly gazes less than 20 sec between shifting of attention; decreased blinking
- 2= Gaze held longer than 20 sec, occasionally shifts attention.
- 3= Fixed gaze, non-reactive.

5. Posturing/ Catalepsy:

Spontaneous maintenance of posture(s), including mundane (e.g., sitting/standing for long periods without reacting).

- 0= Absent
- 1= Less than one minute.
- 2= Greater than one minute, less than 15 minutes.
- 3= Bizarre posture, or mundane maintained more than 15 min.

6. Grimacing:

Maintenance of odd facial expressions.

- 0= Absent
- 1= Less than 10 sec.
- 2= Less than 1 min.
- 3= Bizarre expression(s) or maintained more than 1 min.

7. Echopraxia/ Echolalia:

Mimicking of examiner's movements/ speech.

- 0= Absent
- 1= Occasional.
- 2= Frequent.
- 3= Constant.

8. Stereotypy:

Repetitive, non-goal-directed motor activity (e.g. finger-play; repeatedly touching, patting or rubbing self); abnormality not inherent in act but in its frequency.

- 0= Absent
- 1= Occasional.
- 2= Frequent.
- 3= Constant.

9. Mannerisms:

Odd, purposeful movements (hopping or walking tiptoe, saluting passersby or b

t.
tr

/

Date:

Examiner:

' State examination

' Interval examination over ____ hr.

12. Negativism:

Apparently motiveless resistance to instructions or attempts to move/examine patient. Contrary behavior, does exact opposite of instruction.

- 0= Absent
- 1= Mild resistance and/or occasionally contrary.
- 2= Moderate resistance and/or frequently contrary.
- 3= Severe resistance and/or continually contrary.

13. Waxy Flexibility:

During repositioning of patient, patient offers initial resistance before allowing himself to be repositioned, similar to that of a bending candle.

- 0= Absent.
- 3= Present.

14. Withdrawal:

Refusal to eat, drink and/or make eye contact.

- 0= Absent.
- 1= Minimal PO intake/ interaction for less than one day.
- 2= Minimal PO intake/ interaction for more than one day.
- 3= No PO intake/interaction for one day or more.

15. Impulsivity:

Patient suddenly engages in inappropriate behavior (e.g. runs down hallway, starts screaming or takes off clothes) without provocation. Afterwards can give no, or only a facile explanation.

- 0= Absent
- 1= Occasional
- 2= Frequent
- 3= Constant or not redirectable

16. Automatic Obedience:

Exaggerated cooperation with examiner's request or spontaneous continuation of movement requested.

- 0= Absent
- 1= Occasional
- 2= Frequent
- 3= Constant

17. Mitgehen:

"Anglepoise lamp" arm raising in response to light pressure of finger, despite instructions to the contrary.

- 0= Absent
- 3= Present

18. Gegenhalten:

Resistance to passive movement which is proportional to strength of the stimulus, appears automatic rather than wilful.

- 0= Absent
- 3= Present

19. Ambitendency:

Patient appears motorically "stuck" in indecisive, hesitant movement.

- 0= Absent
- 3= Present

20. Grasp Reflex:

Per neurological exam.

- 0= Absent
- aß= Present M ki p sm â

21. Perseverationf= ° è

Repeatedly returns to same topic or persists with movement.

- 0= Absent
- 3= Present

22. Combativeness: Time:

Usually in an undirected manner, with no, or only a facile explanation afterwards.

- 0= Absent
- 1= Occasionally strikes out, low potential for injury
- 2= Frequently strikes out, moderate potential for injury
- 3= Serious danger to others

23. Autonomic Abnormality:

Circle: temperature, BP, pulse, respiratory rate, diaphoresis.

- 0= Absent
- 1= Abnormality of one parameter [exclude pre-existing hypertension]
- 2= Abnormality of 2 parameters
- 3= Abnormality of 3 or greater parameter